

Composition of Nb-Ta-Ti-Sn-W oxide minerals: indicators of magmatic to hydrothermal evolution of the Cínovec granite intrusion and Sn-W deposit (Czech Republic)

Štěpán Chládek¹, Karel Breiter², Pavel Uher¹

¹ Comenius University, Department of Mineralogy and Petrology; Ilkovičova 6, 842 15 Bratislava, Slovakia; e-mail: st.chladek@seznam.cz

² Institute of Geology of the CAS; Rozvojová 269, 165 00 Praha, Czech Republic; e-mail: breiter@gli.cas.cz

© 2016 Authors. This is an open access publication, which can be used, distributed and reproduced in any medium according to the Creative Commons CC-BY 4.0 License requiring that the original work has been properly cited.

The Cínovec (Zinnwald) Sn-W ore deposit is genetically linked to intrusion of late Variscan, highly fractionated granite which expresses the latest evolutionary stage of a volcano-plutonic system of the Teplíce caldera. Whole intrusion is relatively highly fractionated and from bottom (~1500 m) to top part of cupola-shaped deposit is obviously following succession from biotite (annite) granodiorite-granite-zinnwaldite granite, with the partly greisenized uppermost part at 300–500 m depth (Štemprok 1965, 1971). In 1961–63 the Czechoslovakian Geological Survey (CGS) drilled a 1596 m deep borehole in the Sn-W-mineralized Cínovec granite cupola (Štemprok 1965, Štemprok & Šulcek 1969). All studied rock types include W- and Sn-bearing minerals (wolframite series, scheelite and cassiterite) and disseminated accessory Nb-Ta-Ti-W-Sn minerals (Štemprok & Šulcek 1969, Štemprok 1989, Johan and Johan 1994) which were obtained from the collection of CGS in Prague and studied by BSE and electron microprobe. They crystallized in following succession: rutile + columbite + cassiterite (biotite granodiorite) → rutile + columbite + W-rich ixiolite + cassiterite + scheelite in zinnwaldite granite. Textural relationships of these Nb-Ta-Ti-Sn-W minerals indicate predominantly their magmatic origin and part of them (e.g., cassiterite and columbite) show minor post-magmatic alteration phenomena like distinctly inhomogeneous mixtures of secondary pyrochlore-group minerals

(“oxykenopyrochlore” and oxycalcipyrochlore). Nb/Ta and Fe/Mn fractionation trends led to characteristic Mn and Ta enrichment from bottom (biotite granite) to uppermost zinnwaldite granite, especially in columbite-group minerals. While Nb/Ta fractionation is limitedly applied, effective Fe/Mn fractionation led to significant Mn – enrichment of late-magmatic phases [columbite-(Mn) and W-rich ixiolite].

Post-magmatic to hydrothermal metasomatic fluids caused partial greisenization of the granites and this stage is represented by latest columbite + scheelite + cassiterite + wolframite assemblage. The last two minerals were objects of extensive mining in the past. Although the hydrothermal system was enriched in F and Li (presence of topaz and zinnwaldite), there are only relatively limited Nb/Ta and Fe/Mn fractionations in post-magmatic columbite. Similarly to primary fractionation, both Nb/Ta and Fe/Mn ones take place and overlap characteristic primary Mn-enrichment. Effective Mn-redistribution is predominantly controlled by crystallization of Mn-dominant wolframite like hübnerite in the hydrothermal stage. Scandium is typical rare element in primary (magmatic) and secondary (hydrothermal) mineral assemblage. While primary Sc-fractionation continues the ongoing Sc-enrichment mostly in columbite to uppermost parts of intrusion, the hydrothermal Sc-redistribution is controlled by

crystallization of main ore mineral – wolframite, which consumed a major part of scandium.

Main substitution mechanisms in rutile-cassiterite-wolframite-columbite assemblage include following heterovalent substitutions: (i) $\text{Ti}_3(\text{Fe},\text{Mn})_{-1}^{2+}(\text{Nb},\text{Ta})_{-2}$, (ii) $\text{Ti}_2\text{Fe}_{-1}^{3+}(\text{Nb},\text{Ta})_{-1}$, (iii) $(\text{Nb},\text{Ta})_4\text{Fe}_{-1}^{2+}\text{W}_{-3}$. Moreover, a part of minor cations can enter via: (iv) $(\text{Fe},\text{Mn})_1^{2+}\text{W}_1(\text{Fe},\text{Sc})_{-1}^{3+}(\text{Nb},\text{Ta})_{-1}$ into wolframite lattice, (v) $\text{W}_1(\text{Ti},\text{Sn})_1(\text{Nb},\text{Ta})_{-2}$, (vi) $(\text{Sc},\text{Fe})_3^{3+}(\text{Fe},\text{Mn})_{-2}^{2+}(\text{Nb},\text{Ta})_{-1}$, and (vii) $\text{W}_2\text{Sc}_1^{3+}(\text{Nb},\text{Ta})_{-3}$ into columbite lattice. Calculated Fe^{3+} can be introduced into rutile lattice predominantly via mechanism (ii), while via (iv) into wolframite lattice and together with Sc^{3+} via (vi) into columbite lattice. The last mechanism results in charge imbalance of A and B positions of columbite lattice entering R^{3+} cations to. The distinctly varying calculated Fe^{3+} values can refer to changing $f\text{O}_2$ during columbite, rutile, W-rich ixiolite and wolframite crystallization.

Therefore, the textural and crystallo-chemical features of studied Nb-Ta-Ti-Sn-W oxide minerals

in the Cínovec granite cupola reflect a complex geochemical development of this granite system and ore mineralization from primary magmatic stage, through late-magmatic to subsolidus conditions, and ending in distinct hydrothermally – metasomatic overprint of pre-existing phases.

REFERENCES

- Johan V. & Johan Z., 1994. Accessory minerals of the Cínovec (Zinnwald) granite cupola, Czech Republic. Part 1: Nb-, Ta- and Ti-bearing oxides. *Mineralogy and Petrology*, 51, 323–343.
- Štemprok M., 1965. Petrografie a vertikální rozsah mineralizace v cínovecké žulové klenbě. *Sborník geologických věd. Řada LG: Ložisková geologie*, 5, 7–106.
- Štemprok M., 1971. Petrochemical features of tin-bearing granites in the Krusne Hory Mts., Czechoslovakia. *Society of Mining Geologists of Japan*, Special Issue, 2, 112–118.
- Štemprok M., 1989. Rare earths elements in the rocks of the Cínovec granite cupola (Czechoslovakia). *Věstník Ústředního ústavu geologického*, 64, 1, 1–15.
- Štemprok M. & Šulcek Z., 1969. Geochemical profile through an ore-bearing lithium granite. *Economic Geology*, 64, 4, 392–404.